

Polytechnic
UNIVERSITY

Brooklyn · Long Island · Westchester

On Cell Complexities in Hyperplane Arrangements

Boris Aronov

Micha Sharir



Department of Computer and Information Science

Technical Report

TR-CIS-2000-08

12/15/2000

On Cell Complexities in Hyperplane Arrangements*

Boris Aronov[†]

Micha Sharir[‡]

December 15, 2000

Abstract

We derive improved bounds on the complexity of many cells in arrangements of hyperplanes in higher dimensions, and use these bounds to obtain a very simple proof of a bound, due to [2], on the sum of squares of cell complexities in such an arrangement.

1 Complexity of Many Cells

The main result of the paper, which improves upon previous bounds given in [2], is:

Theorem 1.1 *The complexity of m distinct cells in an arrangement of n hyperplanes in d dimensions, for $d \geq 4$, is $O(m^{1/2}n^{d/2} \log^{(\lfloor d/2 \rfloor - 2)/2} n)$ with the implied constant of proportionality depending on d .*

Proof: The proof proceeds by induction on d . The base case $d = 4$ depends on a sharper bound that is known for $d = 3$ and will be cited below.

Let \mathbf{H} be a collection of n hyperplanes in d -space. We will assume that the planes are in *general position*, meaning that any k planes meet in a $d - k$ -flat, if $k = 1; \dots; d$, and not at all if $k > d$. It is not difficult to see that worst-case cell complexity can always be achieved by planes in general position. Let \mathbf{P} be a set of m points, not lying on any hyperplane. Denote by $\mathbf{K}_j^{(d)}(\mathbf{P}; \mathbf{H})$ the number of j -faces bounding the cells of $\mathcal{A}(\mathbf{H})$ that contain points of \mathbf{P} . We will mainly be concerned with the case $j = \lfloor d/2 \rfloor$, because, as follows from the Dehn-Sommerville relations (see, e.g., [3]), the total number of faces, of all dimensions, of a cell (which is a simple d -polytope) is at most proportional to the number of its $\lfloor d/2 \rfloor$ -faces. We denote by $\mathbf{K}_j^{(d)}(m; n)$ the maximum of $\mathbf{K}_j^{(d)}(\mathbf{P}; \mathbf{H})$ over all sets $\mathbf{P}; \mathbf{H}$ as above.

*Work on this paper has been supported by a grant from the U.S.-Israeli Binational Science Foundation. Work by Boris Aronov was also supported by NSF Grant CCR-99-72568. Work by Micha Sharir was also supported by NSF Grant CCR-97-32101, by a grant from the Israel Science Fund (for a Center of Excellence in Geometric Computing), by the ESPRIT IV LTR project No. 21957 (CGAL), and by the Hermann Minkowski-MINERVA Center for Geometry at Tel Aviv University. Part of the work by Boris Aronov on the paper was done when he visited Tel Aviv University in May 2000.

[†]Department of Computer and Information Science, Polytechnic University, Brooklyn, NY 11201-3840, USA. E-mail: aronov@ziggy.poly.edu

[‡]School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel; and Courant Institute of Mathematical Sciences, New York University, New York, NY 10012, USA. E-mail: sharir@math.tau.ac.il

We now derive a recurrence for $\mathbf{K}_j^{(d)}(\mathbf{m}; \mathbf{n})$. Pick $\mathbf{h} \in \mathbf{H}$, remove it and add it back. Consider the \mathbf{j} -faces that are not contained in \mathbf{h} and bound cells of the arrangement that contain points of \mathbf{P} . This number can increase when \mathbf{h} is added to $\mathcal{A}(\mathbf{H} \setminus \{\mathbf{h}\})$, only when \mathbf{h} splits a cell \mathbf{c} containing points of \mathbf{P} into two subcells, each containing points of \mathbf{P} . In this case, the local increase in the number of \mathbf{j} -faces under consideration is equal to the number of $(\mathbf{j} - 1)$ -faces of the $(\mathbf{d} - 1)$ -face $\mathbf{c} \cap \mathbf{h}$ of $\mathcal{A}(\mathbf{H})$. Denote by $\mathbf{H}=\mathbf{h}$ the set $\{\mathbf{h} \cap \mathbf{h}' \mid \mathbf{h}' \in \mathbf{H} \setminus \{\mathbf{h}\}\}$ of $(\mathbf{d} - 2)$ -hyperplanes within \mathbf{h} . Then the total increase in the number of \mathbf{j} -faces under consideration that is caused by the re-insertion of \mathbf{h} is equal to the number of $(\mathbf{j} - 1)$ -faces in the ‘splitting cells’ of the $(\mathbf{d} - 1)$ -dimensional arrangement $\mathcal{A}(\mathbf{H}=\mathbf{h})$. If the number of cell splittings caused by the re-insertion of \mathbf{h} is \mathbf{m}_h , then the number of \mathbf{j} -faces counted in $\mathbf{K}_j^{(d)}(\mathbf{P}; \mathbf{H})$ and not contained in \mathbf{h} is at most $\mathbf{K}_j^{(d)}(\mathbf{P}_h; \mathbf{H} \setminus \{\mathbf{h}\}) + \mathbf{K}_{j-1}^{(d-1)}(\mathbf{m}_h; \mathbf{n} - 1)$, where \mathbf{P}_h is a subset of \mathbf{P} obtained by removing \mathbf{m}_h points from the cells that got merged when \mathbf{h} was removed. Repeating this analysis for all $\mathbf{h} \in \mathbf{H}$, summing the respective bounds, and taking the maximum over $\mathbf{P}; \mathbf{H}$, we obtain

$$(\mathbf{n} - \mathbf{d} + \mathbf{j})\mathbf{K}_j^{(d)}(\mathbf{m}; \mathbf{n}) \leq \sum_{h \in H} \left(\mathbf{K}_j^{(d)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{K}_{j-1}^{(d-1)}(\mathbf{m}_h; \mathbf{n} - 1) \right); \quad (1)$$

where the factor $\mathbf{n} - \mathbf{d} + \mathbf{j}$ comes from the observation that a \mathbf{j} -face appears in the count for every $\mathbf{h} \in \mathbf{H}$, except for the $\mathbf{d} - \mathbf{j}$ hyperplanes containing it.

The case $\mathbf{d} = 4$. We start with the base case $\mathbf{d} = 4$ (and $\mathbf{j} = 2$). The equation (1) becomes

$$(\mathbf{n} - 2)\mathbf{K}_2^{(4)}(\mathbf{m}; \mathbf{n}) \leq \sum_{h \in H} \left(\mathbf{K}_2^{(4)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{K}_1^{(3)}(\mathbf{m}_h; \mathbf{n} - 1) \right); \quad (2)$$

By the result of [1], we have

$$\mathbf{K}_1^{(3)}(\mathbf{m}; \mathbf{n}) = \begin{cases} \Theta(\mathbf{m}^{2/3}\mathbf{n}) & \text{for } \mathbf{m} \geq \mathbf{n}^{3/2} \\ \Theta(\mathbf{n}^2) & \text{for } \mathbf{n} \leq \mathbf{m} \leq \mathbf{n}^{3/2} \\ \Theta(\mathbf{m}\mathbf{n}) & \text{for } \mathbf{m} \leq \mathbf{n}: \end{cases} \quad (3)$$

Divide (2) by $\mathbf{n}(\mathbf{n} - 1)(\mathbf{n} - 2)$, and put $\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n}) = \mathbf{K}_2^{(4)}(\mathbf{m}; \mathbf{n})/(\mathbf{n}(\mathbf{n} - 1))$, to obtain

$$\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n}) \leq \frac{1}{\mathbf{n}} \sum_{h \in H} \mathbf{F}_2^{(4)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathcal{O} \left(\frac{1}{\mathbf{n}} \sum_{h \in H} \frac{\mathbf{K}_1^{(3)}(\mathbf{m}_h; \mathbf{n} - 1)}{\mathbf{n}^2} \right); \quad (4)$$

We now unwind the recurrence in (4) all the way down to $\mathbf{n}_0 = \mathbf{m}^{1/4}$ remaining hyperplanes. We obtain a recurrence tree \mathbf{T} . The \mathbf{j} -th level of \mathbf{T} is the collection of all nodes whose corresponding substructure involves \mathbf{j} hyperplanes of \mathbf{H} ; thus the root of \mathbf{T} is at level \mathbf{n} (it represents the whole set \mathbf{H}) and the leaves are at level \mathbf{n}_0 . Let π be a path in \mathbf{T} , let $\mathbf{v}_j(\pi)$ denote the node of π at level \mathbf{j} , and let $\mathbf{h}_j(\pi)$ denote the hyperplane removed and reinserted at $\mathbf{v}_j(\pi)$, for $\mathbf{j} = \mathbf{n}; \mathbf{n} - 1; \dots; \mathbf{n}_0 + 1$; in other words, $\mathbf{h}_j(\pi)$ is the hyperplane that represents the edge of π between $\mathbf{v}_j(\pi)$ (parent node) and $\mathbf{v}_{j-1}(\pi)$ (child node). It is easily verified that the unwound recurrence can be rewritten as

$$\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n}) \leq \frac{\mathbf{n}_0!}{\mathbf{n}!} \sum_{\pi} \left[\mathbf{F}_2^{(4)}(\mathbf{m}^*(\pi); \mathbf{n}_0) + \mathcal{O} \left(\sum_{j=\mathbf{n}_0+1}^{\mathbf{n}} \frac{\mathbf{K}_1^{(3)}(\mathbf{m}_j(\pi); \mathbf{j} - 1)}{\mathbf{j}^2} \right) \right]; \quad (5)$$

where γ ranges over all paths in \mathbf{T} , and where $\mathbf{m}_j(\gamma)$ is the number of points removed from the current subset of \mathbf{P} when $\mathbf{h}_j(\gamma)$ is removed from the subset of \mathbf{H} associated with $\mathbf{v}_j(\gamma)$; the number of points remaining in \mathbf{P} after all these removals is denoted by $\mathbf{m}^*(\gamma)$, and we have $\mathbf{m}^*(\gamma) + \sum_{j=n_0+1}^n \mathbf{m}_j(\gamma) = \mathbf{m}$. In other words, $\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n})$ is the average, over all paths of \mathbf{T} , of the path-dependent expression in the brackets in (5). Denote this expression by $\mathbf{E}(\gamma) = \mathbf{F}_2^{(4)}(\mathbf{m}^*(\gamma); \mathbf{n}_0) + \mathbf{O}(\sum_{j=n_0+1}^n \mathbf{E}_j(\gamma))$, where $\mathbf{E}_j(\gamma) = \mathbf{K}_1^{(3)}(\mathbf{m}_j(\gamma); \mathbf{j} - 1) \mathbf{j}^2$.

We fix a path γ in \mathbf{T} , and estimate $\mathbf{E}(\gamma)$. First we have

$$\mathbf{F}_2^{(4)}(\mathbf{m}^*(\gamma); \mathbf{n}_0) = \mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{m}^{1/4}) = \frac{\mathbf{K}_2^{(4)}(\mathbf{m}; \mathbf{m}^{1/4})}{\mathbf{m}^{1/4}(\mathbf{m}^{1/4} - 1)} = \mathbf{O}\left(\frac{\mathbf{O}(\mathbf{m})}{\mathbf{m}^{1/2}}\right) = \mathbf{O}(\mathbf{m}^{1/2});$$

where we have used the fact that an arrangement of $\mathbf{m}^{1/4}$ hyperplanes has $\mathbf{O}(\mathbf{m})$ cells and total complexity $\mathbf{O}(\mathbf{m})$. Partition the nodes of \mathbf{T} into three subsets:

$$\begin{aligned} \mathbf{J}_1 &= \{\mathbf{j} \mid \mathbf{m}_j(\gamma) > (\mathbf{j} - 1)^{3/2}\} \\ \mathbf{J}_2 &= \{\mathbf{j} \mid \mathbf{j} - 1 < \mathbf{m}_j(\gamma) \leq (\mathbf{j} - 1)^{3/2}\} \\ \mathbf{J}_3 &= \{\mathbf{j} \mid \mathbf{m}_j(\gamma) \leq \mathbf{j} - 1\}: \end{aligned}$$

Using (3) and Hölder's inequality, we obtain

$$\begin{aligned} \sum_{j \in \mathbf{J}_1} \mathbf{E}_j(\gamma) &= \mathbf{O}\left(\sum_{j \in \mathbf{J}_1} \frac{\mathbf{m}_j(\gamma)^{2/3}}{\mathbf{j}}\right) \\ &= \mathbf{O}\left[\left(\sum_{j \in \mathbf{J}_1} \mathbf{m}_j(\gamma)\right)^{2/3} \left(\sum_{j > n_0} \frac{1}{\mathbf{j}^3}\right)^{1/3}\right] \\ &= \mathbf{O}\left(\frac{\mathbf{m}^{2/3}}{\mathbf{n}_0^{2/3}}\right) = \mathbf{O}(\mathbf{m}^{1/2}): \end{aligned}$$

Next we have

$$\sum_{j \in \mathbf{J}_3} \mathbf{E}_j(\gamma) = \mathbf{O}\left(\sum_{j \in \mathbf{J}_3} \frac{\mathbf{m}_j(\gamma)}{\mathbf{j}}\right) = \mathbf{O}\left(\sum_{\substack{j \in \mathbf{J}_3 \\ j \leq \mathbf{m}^{1/2}}} \frac{\mathbf{m}_j(\gamma)}{\mathbf{j}} + \sum_{\substack{j \in \mathbf{J}_3 \\ j > \mathbf{m}^{1/2}}} \frac{\mathbf{m}_j(\gamma)}{\mathbf{j}}\right):$$

In the first sum, we use the fact that $\mathbf{m}_j(\gamma) < \mathbf{j}$ to conclude that the sum is $\mathbf{O}(\mathbf{m}^{1/2})$. As for the second sum, we have

$$\sum_{\substack{j \in \mathbf{J}_3 \\ j > \mathbf{m}^{1/2}}} \frac{\mathbf{m}_j(\gamma)}{\mathbf{j}} < \frac{1}{\mathbf{m}^{1/2}} \sum_{\substack{j \in \mathbf{J}_3 \\ j > \mathbf{m}^{1/2}}} \mathbf{m}_j(\gamma) \leq \frac{1}{\mathbf{m}^{1/2}} \cdot \mathbf{m} = \mathbf{m}^{1/2}:$$

Finally, we have

$$\sum_{j \in \mathbf{J}_2} \mathbf{E}_j(\gamma) = \mathbf{O}\left(\sum_{j \in \mathbf{J}_2} 1\right) = \mathbf{O}\left(\sum_{\substack{j \in \mathbf{J}_2 \\ j \leq \mathbf{m}^{1/2}}} 1 + \sum_{\substack{j \in \mathbf{J}_2 \\ j > \mathbf{m}^{1/2}}} 1\right):$$

The first subsum is at most $\mathbf{m}^{1/2}$, while the second is at most

$$\sum_{m_j \geq m^{1/2}} 1 = \frac{\mathbf{m}}{m^{1/2}} = m^{1/2}:$$

To summarize, we have shown that $\mathbf{E}(\cdot) = \mathbf{O}(m^{1/2})$ for each path in \mathbf{T} . Since $\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n})$ is the average of these expressions, we conclude that $\mathbf{F}_2^{(4)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(m^{1/2})$, and hence $\mathbf{K}_2^{(4)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(m^{1/2} \mathbf{n}^2)$. This establishes the base case $\mathbf{d} = 4$, since the Dehn-Sommerville relations imply that $\mathbf{K}_j^{(4)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(\mathbf{K}_2^{(4)}(\mathbf{m}; \mathbf{n}))$, for $\mathbf{j} = 0; 1; 3$, as already mentioned.

The case of odd \mathbf{d} . Next assume that $\mathbf{d} > 4$ is odd, say $\mathbf{d} = 2\mathbf{q} + 1$. In this case, we focus on $\mathbf{j} = \lceil \mathbf{d}/2 \rceil = \mathbf{q} + 1$ and (1) becomes

$$(\mathbf{n} - \mathbf{q})\mathbf{K}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) \leq \sum_{h \in H} \left(\mathbf{K}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{K}_{\mathbf{q}}^{(2\mathbf{q})}(\mathbf{m}_h; \mathbf{n} - 1) \right): \quad (6)$$

By the induction hypothesis, we have

$$\mathbf{K}_{\mathbf{q}}^{(2\mathbf{q})}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(m^{1/2} \mathbf{n}^{\mathbf{q}} \log^{(q-2)/2} \mathbf{n}):$$

We substitute this bound in (6), divide it by $\mathbf{n}(\mathbf{n} - 1) \cdots (\mathbf{n} - \mathbf{q})$, and put $\mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) = \mathbf{K}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) / (\mathbf{n}(\mathbf{n} - 1) \cdots (\mathbf{n} - \mathbf{q} + 1))$, to obtain

$$\mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) \leq \frac{1}{\mathbf{n}} \sum_{h \in H} \mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{O} \left(\frac{1}{\mathbf{n}} \sum_{h \in H} m_h^{1/2} \log^{(q-2)/2} \mathbf{n} \right): \quad (7)$$

We now unwind the recurrence in (7) until only one hyperplane remains. We obtain a recurrence tree \mathbf{T} , and continue to use the same notations as in the case $\mathbf{d} = 4$. It is easily verified that the unwound recurrence can be rewritten as

$$\mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) \leq \frac{1}{\mathbf{n}!} \sum_{\pi} \left[\sum_{j=1}^n \mathbf{O}(m_j(\cdot)^{1/2} \log^{(q-2)/2} \mathbf{j}) \right]; \quad (8)$$

where π ranges over all paths in \mathbf{T} . In other words, as above, $\mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n})$ is the average, over all paths of \mathbf{T} , of the path-dependent expression in the brackets in (8). By the Cauchy-Schwarz inequality, we have

$$\sum_{j=1}^n m_j(\cdot)^{1/2} \leq \left(\sum_{j=1}^n m_j(\cdot) \right)^{1/2} \mathbf{n}^{1/2} \leq m^{1/2} \mathbf{n}^{1/2}:$$

Hence $\mathbf{F}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(m^{1/2} \mathbf{n}^{1/2} \log^{(q-2)/2} \mathbf{n})$, and thus

$$\mathbf{K}_{\mathbf{q}+1}^{(2\mathbf{q}+1)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(m^{1/2} \mathbf{n}^{\mathbf{q}+1/2} \log^{(q-2)/2} \mathbf{n});$$

which is the asserted bound for $\mathbf{d} = 2\mathbf{q} + 1$.

The case of even \mathbf{d} . Finally consider the case where \mathbf{d} is even, say $\mathbf{d} = 2\mathbf{q} > 4$. Here we take $\mathbf{j} = \lceil \mathbf{d}/2 \rceil = \mathbf{q}$. In this case, (1) becomes

$$(\mathbf{n} - \mathbf{q})\mathbf{K}_q^{(2q)}(\mathbf{m}; \mathbf{n}) \leq \sum_{h \in H} \left(\mathbf{K}_q^{(2q)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{K}_{q-1}^{(2q-1)}(\mathbf{m}_h; \mathbf{n} - 1) \right); \quad (9)$$

As noted above, it follows from the Dehn-Sommerville relations that

$$\mathbf{K}_{q-1}^{(2q-1)}(\mathbf{m}_h; \mathbf{n} - 1) = \mathbf{O}(\mathbf{K}_q^{(2q-1)}(\mathbf{m}_h; \mathbf{n} - 1));$$

which allows us to rewrite (9) as

$$(\mathbf{n} - \mathbf{q})\mathbf{K}_q^{(2q)}(\mathbf{m}; \mathbf{n}) \leq \sum_{h \in H} \left(\mathbf{K}_q^{(2q)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{O}(\mathbf{K}_q^{(2q-1)}(\mathbf{m}_h; \mathbf{n} - 1)) \right); \quad (10)$$

By the induction hypothesis, we have

$$\mathbf{K}_q^{(2q-1)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(\mathbf{m}^{1/2} \mathbf{n}^{q-1/2} \log^{(q-3)/2} \mathbf{n});$$

We substitute this bound in (10), divide it by $\mathbf{n}(\mathbf{n} - 1) \cdots (\mathbf{n} - \mathbf{q})$, and put $\mathbf{F}_q^{(2q)}(\mathbf{m}; \mathbf{n}) = \mathbf{K}_q^{(2q)}(\mathbf{m}; \mathbf{n}) / (\mathbf{n}(\mathbf{n} - 1) \cdots (\mathbf{n} - \mathbf{q} + 1))$, to obtain

$$\begin{aligned} \mathbf{F}_q^{(2q)}(\mathbf{m}; \mathbf{n}) &\leq \frac{1}{\mathbf{n}} \sum_{h \in H} \mathbf{F}_q^{(2q)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{O} \left(\frac{1}{\mathbf{n}} \sum_{h \in H} \frac{\mathbf{K}_q^{(2q-1)}(\mathbf{m}_h; \mathbf{n} - 1)}{\mathbf{n}^q} \right) = \\ &\frac{1}{\mathbf{n}} \sum_{h \in H} \mathbf{F}_q^{(2q)}(\mathbf{m} - \mathbf{m}_h; \mathbf{n} - 1) + \mathbf{O} \left(\frac{1}{\mathbf{n}} \sum_{h \in H} \frac{\mathbf{m}_h^{1/2}}{\mathbf{n}^{1/2}} \log^{(q-3)/2} \mathbf{n} \right); \end{aligned} \quad (11)$$

We now unwind the recurrence in (11) until only one hyperplane remains. We obtain a recurrence tree \mathbf{T} , and, as above, rewrite the unwound recurrence as

$$\mathbf{F}_q^{(2q)}(\mathbf{m}; \mathbf{n}) \leq \frac{1}{\mathbf{n}!} \sum_{\pi} \left[\sum_{j=1}^n \mathbf{O} \left(\frac{\mathbf{m}_j(\cdot)^{1/2}}{\mathbf{j}^{1/2}} \log^{(q-3)/2} \mathbf{j} \right) \right]; \quad (12)$$

where π ranges over all paths in \mathbf{T} . In other words, as above, $\mathbf{F}_q^{(2q)}(\mathbf{m}; \mathbf{n})$ is the average, over all paths of \mathbf{T} , of the path-dependent expression in the brackets in (12). By the Cauchy-Schwarz inequality, we have

$$\sum_{j=1}^n \frac{\mathbf{m}_j(\cdot)^{1/2}}{\mathbf{j}^{1/2}} \leq \left(\sum_{j=1}^n \mathbf{m}_j(\cdot) \right)^{1/2} \left(\sum_{j=1}^n \frac{1}{\mathbf{j}} \right)^{1/2} = \mathbf{O}(\mathbf{m}^{1/2} \log^{1/2} \mathbf{n});$$

Hence $\mathbf{F}_q^{(2q)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(\mathbf{m}^{1/2} \log^{(q-2)/2} \mathbf{n})$, and thus $\mathbf{K}_q^{(2q)}(\mathbf{m}; \mathbf{n}) = \mathbf{O}(\mathbf{m}^{1/2} \mathbf{n}^q \log^{(q-2)/2} \mathbf{n})$, which is the asserted bound for $\mathbf{d} = 2\mathbf{q}$.

This completes the proof of the theorem. \square

Remarks: (1) The bounds in the theorem are new, and improve, by a polylogarithmic factor, previous upper bounds given in [2].

(2) In 4 dimensions the bound is $\mathbf{O}(\mathbf{m}^{1/2}\mathbf{n}^2)$. We do not know whether this bound is tight for the whole range of \mathbf{m} . It is clearly tight for $\mathbf{m} = \Theta(1)$ and for $\mathbf{m} = \Theta(\mathbf{n}^4)$. It is also tight for $\mathbf{m} = \Theta(\mathbf{n}^2)$. This has been noted in [2, Theorem 3.3(b)]. For the sake of completeness, here is a sketch of the construction. Take two orthogonal planes $\mathbf{p}; \mathbf{p}'$ in 4-space. Construct in \mathbf{p} an arbitrary arrangement of $\mathbf{n}=2$ lines in general position, and construct in \mathbf{p}' an arrangement of $\mathbf{n}=2$ lines that has a cell \mathbf{c} so that all lines appear on its boundary. Now extend each of these \mathbf{n} lines to a hyperplane in 4-space by taking its Cartesian product with the complementary plane. The cells under consideration in the resulting 4-dimensional arrangement are the Cartesian products of each cell of the arrangement in \mathbf{p} with \mathbf{c} . We obtain a collection of $\mathbf{m} = \Theta(\mathbf{n}^2)$ cells whose overall complexity is $\Theta(\mathbf{n}^2 \cdot \mathbf{n}) = \Theta(\mathbf{n}^3) = \Theta(\mathbf{m}^{1/2}\mathbf{n}^2)$.

(3) The method of proof employed above can also be used to derive the known bound of $\mathbf{O}(\mathbf{m}^{2/3}\mathbf{n}^{d/3} + \mathbf{n}^{d-1})$, $\mathbf{d} \geq 4$, on $\mathbf{K}_{d-1}^{(d)}(\mathbf{m}; \mathbf{n})$, from the corresponding bound in three dimensions. We omit the details.

2 Sum of Squares of Cell Complexities in Hyperplane Arrangements

We next apply Theorem 1.1 to obtain a simple proof of the following result, originally established in [2].

Theorem 2.1 *The sum of squares of the cell complexities in an arrangement of \mathbf{n} hyperplanes in \mathbf{d} dimensions, for $\mathbf{d} \geq 4$, is $\mathbf{O}(\mathbf{n}^d \log^{\lfloor d/2 \rfloor - 1} \mathbf{n})$.*

Proof: Let \mathbf{H} be a set of \mathbf{n} hyperplanes in \mathbf{d} -space, and let $|\mathbf{C}|$ denote the combinatorial complexity (number of faces of all dimensions) of the cell \mathbf{C} in $\mathcal{A}(\mathbf{H})$. We wish to bound the quantity $\Sigma(\mathbf{H}) = \sum_{\mathbf{C}} |\mathbf{C}|^2$.

Remarks: (1) Lemma 3.4 of [2] provides an alternative derivation of this bound from the many-cell bound of Theorem 1.1.

(2) This proof shows that for any $\beta < 2$ we have

$$\sum_C |\mathbf{C}|^\beta = \mathbf{O}(n^d \log^{\lfloor d/2 \rfloor - 2} n):$$

This improves the bound of Theorem 2.1, and, for the cases $\mathbf{d} = 4$ and $\mathbf{d} = 5$, settles in the affirmative a conjecture in [2].

References

- [1] P.K. Agarwal and B. Aronov, Counting facets and incidences, *Discrete Comput. Geom.* 7 (1992), 359–369.
- [2] B. Aronov, J. Matoušek and M. Sharir, On the sum of squares of cell complexities in hyperplane arrangements, *J. Combin. Theory, Ser. A.* 65 (1994), 311–321.
- [3] H. Edelsbrunner, *Algorithms in Combinatorial Geometry*, Springer-Verlag, Heidelberg, 1987.